

KEK-TH-516

April 1997

Mass hierarchy from compositeness hierarchy in supersymmetric gauge theory

M. Hayakawa

*Division of Theoretical Physics, KEK,**Tsukuba, Ibaraki 305, Japan*

(February 1, 2008)

Abstract

The possibility that the mass hierarchy is intimately associated with the compositeness level of the matters is proposed in supersymmetric gauge theory. This implies, for instance, that the preons constituting top quark consists of the “prepreons” binded by the same gauge force making the charm quark out of another preons. The exemplifying toy model illustrates how the hierarchy in the yukawa coupling constants in the up-quark sector is generated from the underlying gauge dynamics. It is also indicated that the incorporation of down-type quarks as elementary objects leads to unpleasant results generically. Thus all the quarks as well as the leptons must also be regarded as composite in the present approach.

Typeset using REVTeX

 JSPS Research Fellow

Electronic address: hayakawa@theory.kek.jp

I. INTRODUCTION

The hierarchic structure observed in the masses among the different generations of quarks and leptons is one of the questions that cannot be addressed to within the standard model. The yukawa sector of the standard model consists of many coupling constants which provide a necessary and sufficient number of parameters to reproduce the masses and mixing of quarks as well as (weak) CP phase at the present experimental stage. In particular there cannot be found any reason why such different orders of numerals appear in the yukawa coupling constants.

In a recent few years various composite models of quarks (and leptons) have been proposed so far [1–3], using the knowledge on the nonperturbative aspects of supersymmetric gauge theory [4]. Those models utilize the dynamics of a series of the asymptotically free gauge theory [5]; confinement without spontaneous breakdown of chiral symmetry. It generates dynamically the nonperturbative terms which accommodate the yukawa interactions of the quarks (and leptons).

In this note we would like to propose another class of composite models to access to the mass hierarchy problem by using the same type of supersymmetric gauge dynamics. That is, the hierarchy of masses may have its origin intimately associated with the *level* of the compositeness of each generation of the matters. It may open up the potential possibility to construct the models possessing more and more predictive power.

The model of Ref. [2] prepares three types of preons, each of which is subject to asymptotically free SU(2) gauge interaction but characterized by the dynamical scales different from each other. Thus, there, the hierarchy of the masses in the up-type quarks, for instance, is traced back to the hierarchy of the scales of the SU(2) gauge dynamics; the repetition of the same structure for confining forces characterized by three different scales.

The matter structure examined here is as follows. For simplicity let us take our attention to the last two generations of the up-type quarks, i.e., charm and top quarks, and consider their respective potential compositeness. The charm quark is assumed to consist of the

preons subject to the confining interactions of the gauge group G_2 . The same force G_2 binds the “prepreons” constituting the preons, which will be binded by asymptotically free G_1 gauge force at lower energy scale to form top quark. Thus as long as we restrict to those two species, the top quarks is considered to be the composite of the composite preons while the charm quark the composite of the preons. To uncover the requisite pattern of hierarchy, Higgs doublet giving the up-type quarks should also be composite with the same level of compositeness as top quark. The same consideration can also be generalized in the case of the three generations.

The next section gives a toy model how such a structure give rise to the hierarchy in the masses of quarks, i.e., the hierarchy in the strength of yukawa coupling, those of which arise from the underlying gauge theory, especially sticking to the two generation. This model itself is far from reality but illustrates the main ingredients of the models based on the idea proposed here.

The last section will be devoted to the discussion and conclusion.

II. EXAMPLE OF PREPREON MODEL

The simplest example of the models realizing the idea described in the previous section is inferred from the use of the infrared dynamics of the supersymmetric $SP(N_c)$ gauge theory with $N_f = N_c + 2$ flavors ¹ of chiral superfields $Q_{\alpha I}$ ² in the fundamental representation [6].

Such a theory gives the confinement of color but does not yield spontaneous breakdown of chiral symmetry $SU(2N_f)$. The massless degrees of freedom appropriate to describe the

¹ The terminology used here obeys that in Ref. [6]. In particular “ N_f flavors” implies that $2N_f$ number [7] of chiral supermultiplets in the fundamental representation (its dimension is $2N_c$) under $SP(N_c)$.

² Here $\alpha = 1, \dots, 2N_c$ denotes the index of the fundamental representation of $SP(N_c)$ and $I = 1, \dots, 2N_f$ are the index of the fundamental the (approximate) global symmetry $SU(2N_f)$.

infrared aspects are the gauge singlet meson chiral superfield $V_{IJ} \sim Q_{\alpha I} Q_{\beta J} \mathcal{J}^{\alpha\beta}$, where $\mathcal{J}^{\alpha\beta}$ is an $\text{SP}(N_c)$ -invariant second-rank antisymmetric tensor, transforming as a second-rank antisymmetric tensor under $\text{SU}(2N_f)$. The gauge theory induces the nonperturbative term to the superpotential which are expressed in terms of this meson field

$$\mathcal{W}_{\text{dyn}} = \text{Pf}_{N_f}(V) \tag{1}$$

$$\equiv \frac{1}{2^{N_f} N_f! \Lambda^{2N_f-3}} \epsilon^{I_1 I_2 \dots I_{2N_f-1} I_{2N_f}} V_{I_1 I_2} \dots V_{I_{2N_f-1} I_{2N_f}}, \tag{2}$$

where Λ is the dynamical scale of $\text{SP}(N_c)$ gauge theory in some renormalization scheme. The aspect of the infrared dynamics of a series of this gauge theory has resemblance to the $\text{SU}(2)$ gauge theory with three flavors (six chiral supermultiplets of spinorial representation), which plays the central role in the models of Ref. [2].

Hereafter we will concentrate on the case of $N_c = 2$ which will turn out to be the minimal choice for the purpose to illustrate how the hierarchy in the yukawa coupling constants arises between the two different generations of up-type quarks (which will be identified as top and charm quarks). The three generation model can be constructed by using $\text{SP}(3) \times \text{SP}(2) \times \text{SU}(2)_T$ as confining gauge forces.

The gauge group of the toy model is $G = \text{SP}(2) \times \text{SU}(2)_T \times G_{\text{ST}}$, where $G_{\text{ST}} = \text{SU}(3)_C \times \text{SU}(2)_L \times \text{U}(1)_Y$ is the gauge group of the standard model. The additional $\text{SU}(2)_T$ is the confining gauge force making the top quarks and the Higgs doubles. Table I lists the content of the particles which are non-singlets under $\text{SP}(2)$ and their charges under G .

The gauge anomaly cancellation with respect to G_{ST} is realized by adding the particles with the charges shown in Table II. Note that the number of each nonzero G_{SM} -charge in $\text{SP}(2)$ non-singlet sector is the same as those of the two generations of Nelson-Strassler model [2]. In order to suppress the baryon number (B) and/or the lepton number violating interactions, the $\text{U}(1)_B$ conservation and Z_2 symmetry are assumed to hold [2] for the charge assignment listed in Table I and II. The fermionic coordinate θ ($\bar{\theta}$) is also charged under

Z_2 in such a way that θ^2 ($\bar{\theta}^2$) has odd parity⁵. Thus Z_2 -charge of each term in the superpotential is required to be odd. The superpotential invariant under $G \times U(1)_B \times Z_2$ for the renormalizable terms containing only the particles in Table I and II is

$$\begin{aligned} \mathcal{W}_{\text{ren-1}} = & \lambda_H \bar{H}_2[\mathbf{h}\mathbf{n}] + \lambda_D \bar{D}_2[\mathbf{d}\mathbf{n}] + \frac{1}{2} \lambda_E E^{(2)}[\mathbf{h}\mathbf{h}] \\ & + \lambda_S \langle [\mathbf{s}\mathbf{s}] \rangle T + \lambda_T T^3 + \sum_{f,g=2}^3 y_L^{fg} \bar{e}_{Rf} l^f \bar{H}_g, \end{aligned} \quad (3)$$

where each $[,]$ implies the contraction of $\text{SP}(2)$ indices through $\mathcal{J}^{\alpha\beta}$, while $\text{SU}(2)_T$ indices are contracted within $<, >$. All the coefficients except for the lepton yukawa coupling constants are $\mathcal{O}(1)$. This should be understood for any tree level superpotential terms hereafter.

For the completion of the toy model, we have to care about the generation of masses for bottom and strange quarks. Unfortunately the present toy model has a disaster on this point. Here we would like to content ourselves with merely presenting a self-contained model and leave it as a future subject.

The renormalizable superpotential each term of which contains at least one of the particles in Table III is as follows;

$$\begin{aligned} \mathcal{W}_{\text{ren-2}} = & M_N \langle \underline{\bar{N}} \underline{N} \rangle + M_P [\bar{\mathbf{P}}\mathbf{P}] + \tau_{\bar{N}} \langle \underline{\bar{N}}[\mathbf{P}\mathbf{s}] \rangle + \tau_N \langle \underline{N}[\bar{\mathbf{P}}\mathbf{s}] \rangle \\ & + \sum_{f=2}^3 \left\{ \tau_H^f \bar{H}_f[\mathbf{P}\mathbf{h}] + \tau_D^f \bar{D}_f[\mathbf{P}\mathbf{d}] + \tau_{\bar{d}_R}^f \bar{d}_{Rf}[\bar{\mathbf{P}}\mathbf{d}] \right\}. \end{aligned} \quad (4)$$

The mass M_P is taken as larger than Λ_{SP} , while M_N should be larger than Λ_T^h , the dynamical scale of $\text{SU}(2)_T$ gauge theory with four spinors. Here it is also assumed to be less than Λ_{SP} .

Below the scale M_P , $\mathbf{P}$ and $\bar{\mathbf{P}}$ decouple from the theory. Integrating out those modes in (4) leads

$$\mathcal{W}_{\text{ren-2}} = M_N \langle \underline{\bar{N}} \underline{N} \rangle$$

⁵ Precisely θ ($\bar{\theta}$) has the Z_4 -charge ω (ω^3) with $\omega = e^{i\pi/2}$ and the imposed symmetry should be called as Z_4 , but this is not important for the present argument. Note that the gaugino has charge +1 so that the soft supersymmetry breaking gaugino mass term is not forbidden by this symmetry.

$$\begin{aligned}
& -\frac{1}{M_P} \left\{ \tau_N \tau_{\bar{N}} \left[\langle \bar{\underline{N}} \underline{\mathbf{s}} \rangle \langle \underline{\mathbf{s}} \underline{N} \rangle \right] + \tau_{\bar{N}} \sum_{f=2}^3 \tau_{\bar{d}_R}^f \bar{d}_{Rf} \langle [\underline{\mathbf{d}} \underline{\mathbf{s}}] \bar{\underline{N}} \rangle \right. \\
& + \tau_N \sum_{f=2}^3 \tau_H^f \bar{H}_f \langle [\underline{\mathbf{h}} \underline{\mathbf{s}}] \underline{N} \rangle + \sum_{f,g=2}^3 \tau_{\bar{d}_R}^f \tau_H^g \bar{d}_{Rf} [\underline{\mathbf{d}} \underline{\mathbf{h}}] \bar{H}_g \left. \right\}, \\
& + \dots,
\end{aligned} \tag{5}$$

where the dots denote the additional terms including D or $\bar{E}$ which are not interested here.

When the energy scale of the system decreases further, SP(2) gauge interaction becomes strong. Then the gauge interactions of G other than SP(2) and the superpotential terms are considered as small compared to the gauge coupling constant of SP(2) above and around the dynamical scale Λ_{SP} so that they can be ignored approximately. Thus the theory is nothing but the supersymmetric SP(2) gauge theory with four flavors of “quarks” $Q_{\alpha I} = (\mathbf{s}, \mathbf{d}, \mathbf{h}, \mathbf{n})_\alpha$ as promised.

The massless composite degrees of freedom consist of the preons $\underline{d}$, $\underline{h}$ and $\underline{n}$ which appear in the models by Nelson and Strassler [2], the singlet S , and the composite chiral supermultiplets $\tilde{V}_{\hat{I}\hat{J}}$ corresponding to the second generation of Nelson-Strassler model with the indices $\hat{I}$ and $\hat{J}$ attached to SU(6) subgroup of the total chiral symmetry SU(8). Explicitly $\tilde{V}_{\hat{I}\hat{J}}$ is expressed as

$$\tilde{V}_{\hat{I}\hat{J}} \sim \Lambda_{\text{SP}} \begin{pmatrix} \bar{c}_R & q_L^{(2)} & -D^{(2)} \\ -q_L^{(2)} & \bar{E}_2 & -H^{(2)} \\ D^{(2)} & H^{(2)} & 0 \end{pmatrix}. \tag{6}$$

There appear the right-handed charm quark $\bar{c}_R$, the left-handed quark doublet $q_L^{(2)}$ of the second generation, and the Higgs doublets $H^{(2)}$, which will not obtain nonzero vacuum expectation value (VEV) as its mass squared will be found of the order Λ_{SP}^2 , much larger than the negative corrections by the large top yukawa coupling of the order $-(y_t m_{\text{susy}}/(4\pi))^2$ with the supersymmetry breaking scale $m_{\text{susy}} \sim \mathcal{O}(1)$ TeV and top yukawa coupling constant y_t of order one.

In contrast to the Nelson-Strassler model the nonperturbatively generated superpotential terms are now the non-renormalizable ones *ab initio*

$$W_{\text{dyn}} = \text{Pf}_2(V), \quad (7)$$

which explicitly give rise to

$$\begin{aligned} \Lambda_{\text{SP}} W_{\text{dyn}} = S \Big\{ & \alpha'_2 \bar{c}_R q_L^{(2)} H^{(2)} + \beta'_2 \bar{c}_R D^{(2)} \bar{E}_2 + \gamma'_2 D^{(2)} q_L^{(2)} q_L^{(2)} \Big\} \\ & + \alpha_2^H \bar{c}_R q_L^{(2)} \langle \underline{h} \underline{n} \rangle + \alpha_2^{t_R} \langle \underline{d} \underline{d} \rangle q_L^{(2)} H^{(2)} + \alpha_2^{q_L} \bar{c}_R \langle \underline{d} \underline{h} \rangle H^{(2)} \\ & + \beta_2^{t_R} \langle \underline{d} \underline{d} \rangle D^{(2)} \bar{E}_2 + \beta_2^D \bar{c}_R \langle \underline{d} \underline{n} \rangle \bar{E}_2 + \beta_2^E \bar{c}_R D^{(2)} \langle \underline{h} \underline{h} \rangle \\ & + \gamma_2^{q_L} \langle \underline{d} \underline{h} \rangle q_L^{(2)} D^{(2)} + \gamma_2^D \langle \underline{d} \underline{n} \rangle q_L^{(2)} q_L^{(2)}, \end{aligned} \quad (8)$$

where α' etc., are $\mathcal{O}(1)$ constants parametrizing the corrections from the perturbative couplings, which depend on the species taking part in each above coupling. The $U(1)_B$ is conserved by the dynamically generated terms since they do not break chiral symmetry $SU(8)$ and the baryon number restricted to $SP(2)$ non-singlet sector is one of the generators of $SU(8)$. The tree level superpotential in (3) now turns into

$$\begin{aligned} \mathcal{W}_{\text{ren-1}} = & \lambda_H \Lambda_{\text{SP}} \bar{H}_2 H^{(2)} + \lambda_D \Lambda_{\text{SP}} \bar{D}_2 D^{(2)} + \lambda_E \Lambda_{\text{SP}} E^{(2)} \bar{E}_2 + \lambda_S \Lambda_{\text{SP}} S T \\ & + \lambda_T T^3 + \sum_{f,g=2}^3 y_L^{fg} \bar{e}_R f l^f \bar{H}_g. \end{aligned} \quad (9)$$

Eq. (5) also becomes

$$\begin{aligned} \mathcal{W}_{\text{ren-2}} = & M_N \langle \bar{N} N \rangle \\ & - \frac{\Lambda_{\text{SP}}}{M_P} \left\{ \sum_{f=2}^3 \left(\tau_{\bar{N}} \tau_{\bar{d}_R}^f \bar{d}_{Rf} \langle \underline{d} \bar{N} \rangle + \tau_N \tau_H^f \bar{H}_f \langle \underline{h} N \rangle \right) + \sum_{f,g=2}^3 \tau_{\bar{d}_R}^f \tau_H^g \bar{d}_{Rf} q_L^{(2)} \bar{H}_g \right\} \\ & + \dots \end{aligned} \quad (10)$$

Now $H^{(2)}$, $\bar{H}_2$, $D^{(2)}$, $\bar{D}_2$, $E^{(2)}$, E_2 , S and T gain masses of order Λ_{SP} and decouple from the effective theory under those thresholds, producing new non-renormalizable operators, but irrelevant for mass generation. Keeping the relevant terms for the quark yukawa couplings explicitly in the sum of (8), (9) and (10) the effective superpotential becomes

$$\begin{aligned} \mathcal{W}_{\text{eff}} = & \alpha_2^H \frac{1}{\Lambda_{\text{SP}}} \bar{c}_R q_L^{(2)} \langle \underline{h} \underline{n} \rangle \\ & - \frac{\Lambda_{\text{SP}}}{M_P} \left\{ \tau_{\bar{N}} \sum_{f=2}^3 \tau_{\bar{d}_R}^f \bar{d}_{Rf} \langle \underline{d} \bar{N} \rangle + \tau_N \tau_H \bar{H} \langle \underline{h} N \rangle + \tau_H \sum_{f=2}^3 \tau_{\bar{d}_R}^f \bar{d}_{Rf} q_L^{(2)} \bar{H} \right\} \\ & + \dots, \end{aligned} \quad (11)$$

by writing $\bar{H}$ for $\bar{H}^{(3)}$.

As the energy further goes down, $\underline{N}$, $\bar{\underline{N}}$ decouples and the theory becomes described by the effective superpotential

$$\begin{aligned} \mathcal{W}_{\text{eff}} = & \alpha_2^H \frac{1}{\Lambda_{\text{SP}}} \bar{c}_R q_L^{(2)} \langle \underline{h} \underline{n} \rangle - \tau_H \frac{\Lambda_{\text{SP}}}{M_P} \sum_{f=2}^3 \tau_{\bar{d}_R}^f \bar{d}_{Rf} q_L^{(2)} \bar{H} \\ & - \frac{\Lambda_{\text{SP}}}{M_P M_N} (\tau_{\bar{N}} \tau_N \tau_H) \sum_{f=2}^3 \tau_{\bar{d}_R}^f \bar{d}_{Rf} \langle \underline{d} \underline{h} \rangle \bar{H} + \dots \end{aligned} \quad (12)$$

Finally the gauge force $\text{SU}(2)_T$ becomes strong at the scale Λ_T enough to confine the composite preons $\underline{d}$, $\underline{h}$ and $\underline{n}$, where Λ_T is now the dynamical scale of $\text{SU}(2)_T$ gauge theory with six spinors. This is the dynamical feature due to the supersymmetric $\text{SU}(2)$ gauge theory with six spinors. The massless composite degrees of freedom are then

$$\begin{pmatrix} \langle \underline{d} \underline{d} \rangle & \langle \underline{d} \underline{h} \rangle & \langle \underline{d} \underline{n} \rangle \\ \langle \underline{h} \underline{d} \rangle & \langle \underline{h} \underline{h} \rangle & \langle \underline{h} \underline{n} \rangle \\ \langle \underline{n} \underline{d} \rangle & \langle \underline{n} \underline{h} \rangle & 0 \end{pmatrix} \sim \Lambda_T \begin{pmatrix} \bar{t}_R & q_L^{(3)} & -D^{(3)} \\ -q_L^{(3)} & \bar{E}_3 & -H \\ D^{(3)} & H & 0 \end{pmatrix}. \quad (13)$$

$\text{SU}(2)_T$ gauge dynamics generates the additional new cubic terms to the superpotential (12) nonperturbatively

$$\begin{aligned} \mathcal{W}_{\text{eff}} = & \alpha \bar{t}_R q_L^{(3)} H + \alpha_2^H \frac{\Lambda_T}{\Lambda_{\text{SP}}} \bar{c}_R q_L^{(2)} H - \tau_H \frac{\Lambda_{\text{SP}}}{M_P} \sum_{f=2}^3 \tau_{\bar{d}_R}^f \bar{d}_{Rf} q_L^{(2)} \bar{H} \\ & - \frac{\Lambda_{\text{SP}} \Lambda_T}{M_P M_N} (\tau_{\bar{N}} \tau_N \tau_H) \sum_{f=2}^3 \tau_{\bar{d}_R}^f \bar{d}_{Rf} q_L^{(3)} \bar{H} + \dots \end{aligned} \quad (14)$$

Eq. (14) shows that the yukawa coupling constant for top quark appears as $\mathcal{O}(1)$ [1,2] while that for charm quark is damped by the factor $\Lambda_T/\Lambda_{\text{SP}}$, the ratio of the confining scales of the two gauge dynamics. Note that the origin of this factor arises from the non-renormalizable nature of $\text{SP}(N_c)$ dynamics and the multi-compositeness of the Higgs doublet acquiring nonzero VEV. However as one linear combination of $\bar{d}_R$ participates in (14), only one down-type quark gains the mass, which is strange quark as the yukawa coupling in matter involves $q_L^{(2)}$. This unpleasant feature in the present example originates from the multi-compositeness of $q_L^{(3)}$ which prevents a renormalizable superpotential term and leaves

a global symmetry in the yukawa-sector. Such a pattern that bottom becomes lighter than strange quark will not be circumvented as long as these quarks are considered as elementary objects since the dimension of the operator containing the third-generation quark doublet $q_L^{(3)}$ necessarily becomes higher than that for $q_L^{(2)}$ when they are written down in terms of their most fundamental objects. Obviously such a feature originates from our present insight on the structure of matters. However compositeness of the right-handed down-type quarks will turn over this invalid relations if confinement producing bottom quark occurs at much higher scale than that making strange quark, or strange quark has deeper substructure than bottom quark.

III. DISCUSSION AND CONCLUSION

Here we seek the possibility that the mass hierarchy of the matters is directly related with the compositeness hierarchy of the matters. The toy model is given to illustrate that the dynamically generated nonrenormalizable terms induces the suppression factor for the light up-type quark yukawa coupling. This factor is the ratio of the two confining scales, differing from that in Ref. [1] using similar $SP(N_c)$ gauge dynamics, in which the yukawa coupling for the light flavors arises from the non-renormalizable terms expected to be induced from Planck scale physics. The compositeness of Higgs doublet H also plays an essential role here.

It was also observed that the down-type quarks must be considered as composite to reach the realistic pattern of their masses once we follow up the present idea. Accepting this fact positively, this requirement naturally suggests that all the matters should be composite if we follow the present approach.

The multi-compositeness of the matter often forbids the renormalizable superpotential at tree level assuring an accidental symmetry. Such an inherent property might become useful to generate desirable hierarchy in the cleverer model, although it yielded some unpleasant results (including μ problem as discussed below) in the toy model here.

The exemplifying model does not generate the mass term for Higgs bosons (so called μ term) as well as the mass terms for $D^{(3)}$, $\bar{D}_3$, $E^{(3)}$ and $\bar{E}_3$ from the renormalizable interactions. The nonrenormalizable operators to induce it can always be put in by hand, for instance, as

$$\frac{1}{M_1 M_2} \bar{H} \langle [\mathbf{hs}][\mathbf{sn}] \rangle, \quad (15)$$

for the Higgs mass terms, but it is rather unattractive as the mass scale entering above must be tuned to give a right order of magnitude as $\Lambda_{\text{SP}}^2 \Lambda_{\text{T}} / (M_1 M_2) \sim \mathcal{O}(1)$ [TeV]. This point also needs further consideration in the construction of the realistic model based on the present approach, which is left over as the future subject.

ACKNOWLEDGMENTS

The author thanks N. Kitazawa for discussing about the supersymmetric gauge theory.

REFERENCES

- [1] M. J. Strassler, Phys. Lett. **B376** (1996) 119.
- [2] A. E. Nelson and M. J. Strassler, UW-PT-96-09, hep-ph/9607362 (1996).
- [3] M. A. Luty and R. N. Mohapatra, UMD-PP-97-43, hep-ph/9611343 (1996).
- [4] N. Seiberg, Phys. Lett. **B318** (1993) 469; Phys. Rev. D **49** (1994) 6857; K. Intriligator, R. G. Leigh and N. Seiberg, Phys. Rev. D **50** (1994) 1092; K. Intriligator and N. Seiberg, Nucl. Phys. Proc. Suppl. **45BC** (1996) 1.
- [5] C. Csáki, M. Schmaltz and W. Skiba, MIT-CTP-2597, BUHEP-96-46, hep-th/9612207 (1996).
- [6] K. Intriligator and P. Pouliot, Phys. Lett. **B353** (1995) 471.
- [7] E. Witten, Phys. Lett. **B117** (1982) 324.

TABLES

TABLE I. SP(2) non-singlet species and their quantum numbers. $U(1)_B$ is the baryon number.

	SP(2)	SU(2) _T	SU(3) _C	SU(2) _L	U(1) _Y	U(1) _B	Z_2
s	4	2	1	1	0	0	+
d	4	1	3	1	$-\frac{1}{3}$	$-\frac{1}{6}$	+
h	4	1	1	2	$\frac{1}{2}$	$\frac{1}{2}$	+
n	4	1	1	1	0	$-\frac{1}{2}$	−

TABLE II. SP(2) singlet species and their charges under symmetry group, in which each of them is introduced in duplicate ($f = 2, 3$), except for T .

	SP(2)	SU(2) _T	SU(3) _C	SU(2) _L	U(1) _Y	U(1) _B	Z_2
$\bar{d}_{Rf}$	1	1	3*	1	$\frac{1}{3}$	$-\frac{1}{3}$	−
$\bar{H}_f$	1	1	1	2	$-\frac{1}{2}$	0	+
E^f	1	1	1	1	−1	−1	−
$\bar{D}_f$	1	1	3*	1	$\frac{1}{3}$	$\frac{2}{3}$	+
$\bar{e}_{Rf}$	1	1	1	1	1	0	+
l^f	1	1	1	2	$-\frac{1}{2}$	0	−
T	1	1	1	1	0	0	−

TABLE III. Additional species incorporated for generating the down-type quark masses.

	SP(2)	SU(2) _T	SU(3) _C	SU(2) _L	U(1) _Y	U(1) _B	Z_2
$\underline{N}$	1	2	1	1	0	$-\frac{1}{2}$	−
$\underline{\bar{N}}$	1	2	1	1	0	$\frac{1}{2}$	+
P	4	1	1	1	0	$-\frac{1}{2}$	−
$\bar{\mathbf{P}}$	4	1	1	1	0	$\frac{1}{2}$	+